

しゅるい 種類	のぞ じき 望ましい時期	かいすう 回数	かんかく 間隔
ポリオ えきたい の (液体を飲みます)	せいご かげつ さい かげつ 生後3ヶ月～1歳6ヶ月ごろ	かい 2回	4～6 週
DPT D:ジフテリア P:百日咳 T:破傷風	き せいご かげつ さい かげつ 1期:生後3ヶ月～2歳6ヶ月ごろ き 2期(DTのみ):11歳	かい 4回 かい 1回	3～8 週 あけて3回 かいめ ねんご 4回目は1年後
ましん ふうしん 麻疹・風疹 (MR)	き さい さい 1期:1歳～2歳 き さい さいみまん しょうがっこうしゅうがくまえ 1ねんかん 2期:5歳～7歳未満で小学校就学前の1年間	かい 2回	
BCG	せいご かげつ かげつ 生後3ヶ月～6ヶ月	かい 1回	

An example of a schedule for regular vaccinations

Type	a desirable period for vaccination	Number of doses	Interval
Poliomyelitis (taken orally as a liquid)	From three months old to one year and half	twice	Four to six weeks
DPT D: diphtheria P: pertussis (whooping cough) T: tetanus	The first period: From three months old to two years and a half The second period (only DT): eleven years old	Four times once	Three times ; The interval between doses is three to eight weeks The Fourth time: at least one year after the third time
Measles & Rubella (MR)	The first period: From one year to two years old The second period: five years old to less than seven years old, which means one year before enrolling in elementary school	twice	
BCG	From three months to six months old	once	